

Immune Evasion Mechanisms of the Zoonotic Protozoan Parasite *Toxoplasma Gondii* in Mammalian Hosts

著者	BANDO Hironori, FUKUDA Yasuhiro, YAMAMOTO Masahiro, KATO Kentaro
journal or publication title	Journal of Integrated Field Science
volume	18
page range	13-17
year	2021-03
URL	http://hdl.handle.net/10097/00131377



Symposium mini review

Immune Evasion Mechanisms of the Zoonotic Protozoan Parasite *Toxoplasma Gondii* in Mammalian Hosts

Hironori BANDO^{1,2}, Yasuhiro FUKUDA¹, Masahiro YAMAMOTO² and Kentaro KATO¹

¹Laboratory of Sustainable Animal Environment, Graduate School of Agricultural Science, Tohoku University, 232-3 Yomogida, Naruko-onsen, Osaki, Miyagi 989-6711, Japan

²Department of Immunoparasitology, Research Institute for Microbial Diseases, Osaka University, 3-1, Yamadaoka, Suita, Osaka, 565-0871, Japan

Keywords

Toxoplasma, IFN- γ , NLRP3, iNOS, TgGRA15

Corresponding Author

Kentaro KATO,
kentaro.kato.c7@tohoku.ac.jp

Abstract

Toxoplasma gondii is a zoonotic protozoan pathogen that causes toxoplasmosis, an infectious disease that affects most mammals, including domestic animals, wild animals, and humans. Toxoplasmosis in domestic animals causes miscarriages or stillbirths, resulting in economic losses and posing a challenge in animal husbandry. *T. gondii* is thus an important pathogen that causes serious animal and public health issues, yet there is still no vaccine or preventative medicine. Therefore, efforts to develop novel treatments for toxoplasmosis and to understand the interaction between the host immune response and the parasite in host cells are essential. We know that interferon- γ (IFN- γ)-induced tryptophan degradation by indole-2,3-dioxygenase (IDO1) plays an important role in the IFN- γ -induced anti-*T. gondii* response. However, little is known about *T. gondii* virulence mechanisms targeting IDO1. Therefore, we focused on the *T. gondii* effector TgGRA15 and analyzed its virulence function and mechanism to antagonize the IDO1-mediated anti-*T. gondii* response. In this study, we demonstrate that inducible nitric oxide synthase is a key host factor for TgGRA15-dependent disruption of the IDO1-dependent anti-*T. gondii* response.

Introduction

Toxoplasma gondii is an obligate intracellular zoonotic protozoan parasite that causes toxoplasmosis in most mammals, including domestic animals, wild animals, and humans (Boothroyd, 2009; Dubey, 2010). The family *Felidae*, which includes domestic cats, is the definitive host of *T. gondii*. The parasite can easily spread infection through the accidental swallowing of food or water contaminated with oocysts. Accordingly, *T. gondii* is prevalent in most areas of the world (Montazeri *et al.*, 2020). Toxoplasmosis in humans and domestic animals can cause congenital disease, miscarriages and stillbirths, leading to not only problems of animal hygiene and public health, but also economic losses to farmers (Stelzer *et al.*, 2019). Yet, no effective vaccine or preventive drug has yet been developed. Recently, along with increased overlap of the living space between humans, domestic animals, and wild animals, the number of cases of toxoplasmosis has been increasing annually. In fact, in 2015, we reported that the

number of *T. gondii* infected-wild animals is increasing in Japan (Bando *et al.*, 2015). Furthermore, *T. gondii* has been ranked among the top five human pathogens that cause life impairment and economic losses in the United States (Batz *et al.*, 2012). Therefore, to develop novel therapeutic methods or medicines against *T. gondii*, basic research on the interaction between *T. gondii* and its host is essential.

The host immune resistance responses to *T. gondii* rely on innate and adaptive immunity (Lee *et al.*, 2015; Ma *et al.*, 2014; MacMicking, 2012). Interferon- γ (IFN- γ), which is produced by CD4⁺ T cells and natural killer cells and stimulates cell-autonomous responses in both immune and non-immune cells, is the most important molecule for anti-*T. gondii* responses (Suzuki *et al.*, 1988). IFN- γ plays a role in the activation of the STAT1 transcription factor and induction of the expression of hundreds of genes (Platanias, 2005). Some studies have shown that IFN- γ -inducible GTPases mediate parasitocidal and parasitostatic responses in mice (Taylor *et al.*, 2007; Zhao *et al.*, 2009; Yamamoto *et al.*, 2012), whereas other recent

studies have reported that these GTPases may not play major roles in IFN- γ dependent anti-*T. gondii* responses in human cells (Ohshima *et al.*, 2015; Fisch *et al.*, 2019). We have shown that IFN- γ stimulates the expression of indoleamine 2,3-dioxygenase (IDO) and has an essential role in the anti-*T. gondii* responses of various human cell types (Bando *et al.*, 2018b). Thus, although IFN- γ has a critical role in the anti-*T. gondii* response of both humans and mice, the IFN- γ -inducible effector mechanisms may differ between these two species.

T. gondii secretes various effector molecules, called rhoptry proteins (ROPs) and dense granule proteins (GRAs), into host cells. These effectors are frequently used to promote parasite growth in host cells (Hakimi *et al.*, 2017; Hunter and Sibley, 2012), and their virulence mechanisms, function, and significance have been extensively researched in mouse models (Behnke *et al.*, 2011; Etheridge *et al.*, 2014; Fentress *et al.*, 2010; Reese *et al.*, 2011; Rosowski *et al.*, 2014; Rosowski and Saeij, 2012; Steinfeldt *et al.*, 2010). The *Toxoplasma* effector TgGRA15, one of the dense granule proteins, is secreted into host cells to activate the host transcription factor NF- κ B in mice (Gov *et al.*, 2013; Jensen *et al.*, 2011; Rosowski *et al.*, 2011), although it should be noted that most virulence factors suppress the host immune responses (Olias *et al.*, 2016; Gay *et al.*, 2016). TgGRA15-deficient *T. gondii* has been shown to promote parasite proliferation *in vivo* in mice (Jensen *et al.*, 2013; Rosowski *et al.*, 2011), meaning that TgGRA15 can support host survival by preventing parasite growth. Thus, the significance of TgGRA15 as a virulence factor remains unclear. In this study, we introduce the virulent mechanism of TgGRA15 targeting the IDO1-dependent anti-*T. gondii* response in human cells.

TgGRA15 promotes *T. gondii* growth when co-cultured in the presence of IFN- γ

The function of TgGRA15 as a virulence factor is unclear; therefore, to explore it in human cells, we generated TgGRA15-deficient (TgGRA15-KO) *T. gondii* by using the CRISPR/Cas9 system. Then, we tested whether TgGRA15 has an important role in the suppression of host immune responses under human cell mono-culture conditions. However, we failed to find any advantageous effect of TgGRA15 on parasite growth in various human cell lines. When *T. gondii* infects its host, the parasite preferentially infects CD11b⁺ cells such as monocytes, and then the infected cells are carried by the bloodstream to various organs (Courret *et al.*, 2006). Several kinds of co-culture models have been established to mimic complex cell-cell interactions by using human tissue or immune cell lines, one of which is the monocyte-hepatocyte co-culture model (Frenkel and Remington, 1980). Because one of the major symptoms of toxoplasmosis is hepatitis, we developed a *T. gondii* infection model using monocyte-hepatocyte co-culture conditions. Human acute monocytic leukemia cell line THP-1 cells were infected with wild-type or TgGRA15-KO parasite, and then both the culture supernatant and infected THP-1 cells were seeded onto human hepatoma cell line Huh7 cells with or without IFN- γ . Interestingly, the parasite numbers under the TgGRA15-KO parasite-infected co-culture condition were significantly reduced compared with the wild-type parasite-infected co-culture condition. These

data indicate that TgGRA15 has an advantageous effect on *T. gondii* growth under human cell co-culture conditions.

NLRP3-dependent IL-1 β secretion from monocytes is essential for the pro-parasitic effect of TgGRA15 in hepatocytes

We next attempt to reveal the mechanisms of the pro-parasitic effect of TgGRA15 under co-culture conditions. First, to test whether TgGRA15 has an effect on monocytes or hepatocytes, the culture supernatants were collected from wild-type and TgGRA15-KO *T. gondii*-infected THP-1 cells, and then both the parasites and THP-1 cells were removed by filtration. The filtered culture supernatants and newly prepared wild-type or TgGRA15-KO parasites were then added to Huh7 cells with IFN- γ . Then the number of parasites in the Huh7 cells was assessed. The presence of TgGRA15 in THP-1 cells, but not Huh7 cells, led to a reduction in parasite number, suggesting that the presence of TgGRA15 in monocytes and their supernatant is essential for the pro-parasitic effect. Therefore, we next focused on the components of the supernatant from the parasite infected-THP-1 cell culture. Previous studies have reported that *T. gondii* infection induces proinflammatory cytokine IL-1 β secretion from THP-1 cells in a TgGRA15-dependent manner (Gov *et al.*, 2013). It has also been reported that IL-1 β production in monocytes is dependent on Caspase-1 and inflammasome activation (Gov *et al.*, 2013; Gov *et al.*, 2017). Therefore, to test whether TgGRA15-dependent Caspase-1 and inflammasome activation are important for IL-1 β secretion from monocytes, we generated NLRP3-deficient (NLRP3-KO) or Caspase-1-deficient (CASP1-KO) THP-1 cells by using CRISPR/Cas9 systems, and then analyzed IL-1 β secretion levels in the culture supernatant. We found that both NLRP3-KO- and CASP1-KO-infected THP-1 cells showed significantly reduced IL-1 β secretion. Then, we examined whether IL-1 β secretion in THP-1 cells is essential for suppressing the IFN- γ -dependent anti-*T. gondii* response under co-culture conditions. We found that the parasite number in both wild-type parasite-infected NLRP3-KO and CASP1-KO THP-1 cells was significantly reduced compared with that of wild-type THP-1 cells. These results indicate that IL-1 β secretion through Caspase-1 and NLRP3 inflammasome activation in THP-1 cells has an important role in the TgGRA15-dependent suppression of the IFN- γ -dependent anti-*T. gondii* response.

The IFN- γ -induced IDO1-dependent anti-*T. gondii* response is downregulated by TgGRA15 in hepatocytes

We previously reported that IDO1-induced tryptophan degradation has an important role in the IFN- γ -dependent anti-*T. gondii* response in various human cell types including hepatocytes (Bando *et al.*, 2018b; Bando *et al.*, 2019) because tryptophan is an essential amino acid for parasite growth. In fact, we found that the IFN- γ -dependent reduction in parasite numbers in IDO1-deficient (IDO1-KO) Huh7 cells was abolished under TgGRA15-KO parasite-infected co-culture conditions. Therefore, we examined whether IL-1 β affects IDO1 expression in Huh7 cells. We found that IL-

IL-1 β and IFN- γ co-stimulation severely inhibited IDO1 mRNA and protein levels in Huh7 cells. Then, to examine whether IL-1 β -dependent impairment of the IFN- γ -dependent anti-*T. gondii* response was IDO1-dependent, we generated MyD88-deficient (MyD88-KO)—MyD88 is essential molecule for the IL-1 receptor signaling pathway (Adachi *et al.*, 1998)—and IL1R1-deficient (IL1R1-KO) Huh7 cells by using CRISPR/Cas9 systems. We found that the pro-parasitic effect of IL-1 β in IDO1-KO, MyD88-KO, and IL1R1-KO Huh7 cells was completely abolished. Then, we compared the protein levels of IDO1 under wild-type *T. gondii*- and TgGRA15-KO parasite-infected and non-infected co-culture conditions. We found that the protein levels of IDO1 were significantly reduced under wild-type parasite-infected conditions compared with non-infected conditions. Importantly, the protein levels of IDO1 under TgGRA15-KO parasite-infected conditions recovered to the same levels as those seen under non-infected conditions. These results indicate that the TgGRA15-induced IL-1 β -dependent downregulation of IDO1 expression is important for the impairment of the IFN- γ -dependent anti-*T. gondii* response in hepatocytes.

iNOS is essential for TgGRA15-dependent inhibition of the IDO1-dependent anti-*T. gondii* response

Nitric oxide (NO) production is known to strongly downregulate IDO activity transcriptionally, translationally, and post-translationally (Thomas *et al.*, 1994). In addition, inducible nitric oxide synthase (iNOS) has been shown to be an important factor for IFN- γ -mediated NO production (Nathan and Xie, 1994). Hence to explain the mechanism of the IL-1 β -dependent IDO1 suppression, we focused on iNOS and NO-dependent downregulation of IDO1 activity in hepatocytes. First, we examined the expression level of iNOS mRNA in Huh7 cells. We found that IFN- γ and IL-1 β co-stimulation enhanced the expression level of iNOS mRNA and strongly induced NO production in Huh7 cells. Then, to examine the role of iNOS, we generated iNOS-deficient (iNOS-KO) Huh7 cells by using the CRISPR/Cas9 system. We found that NO was not produced from iNOS-KO Huh7 cells upon IFN- γ and IL-1 β co-stimulation and that IL-1 β -dependent reduction of IDO1 protein levels did not occur in iNOS-KO Huh7 cells. Furthermore, the IL-1 β -dependent pro-parasitic effect was completely abolished in the iNOS-KO Huh7 cells under mono-culture conditions, suggesting that IL-1 β induces iNOS expression to inhibit the IDO1-dependent anti-*T. gondii* response. Then, we tested whether this mechanism occurs under co-culture conditions. We found that NO production and the reduction of IDO1 was not observed in iNOS-KO Huh7 cells co-cultured with wild-type parasite-infected wild-type THP-1 cells. Moreover, the TgGRA15-dependent pro-parasitic effect was abolished in iNOS-KO Huh7 cells under these co-culture conditions. Finally, we confirmed the GRA15-dependent virulence mechanism in primary human cells. Taken together, our results indicate that iNOS is an essential host factor for the TgGRA15-dependent virulence mechanism under monocyte-hepatocyte co-culture conditions.

Conclusion

In summary, here we showed that IL-1 β is produced from monocytes in a *Toxoplasma* effector TgGRA15- and host NLRP3 inflammasome-dependent manner. We further showed that iNOS has an essential role in the *Toxoplasma* TgGRA15-dependent inhibition of the IDO1-induced anti-*T. gondii* response in human cells (Bando *et al.*, 2018a). Although immune responses in humans and domestic animals are not identical (Guzman and Montoya, 2018), the tryptophan-degrading enzyme IDO and nitric oxide synthases NOS have been found in most mammalian species (Yao *et al.*, 2011). Hence, the TgGRA15-dependent virulence mechanism may contribute to *T. gondii* infection in not only humans but also domestic animals. Studies to examine whether the TgGRA15-dependent virulence mechanism has an important role in *T. gondii* infection of domestic animals, and to identify chemical compounds that block iNOS expression or NO production could contribute to the development of novel anti-toxoplasmosis therapies for humans and domestic animals.

Acknowledgments

We thank the staff at the Laboratory of Immunoparasitology, Research Institute for Microbial Diseases, Osaka University and the Laboratory of Sustainable Animal Environment, Graduate School of Agricultural Science, Tohoku University. This study was supported by a Research Program on Emerging and Reemerging Infectious Diseases (17fk0108120h0001) and a fund for the Promotion of Joint International Research (Fostering Joint International Research (B)) (19KK0242) from the Japan Society for the Promotion of Science (JSPS) and the Japanese Initiative for Progress of Research on Infectious Diseases for Global Epidemic (17fm0208018h0001) from the Agency for Medical Research and Development (AMED), by a Grant-in-Aid for Scientific Research on Innovative Areas (17K15677 and 19K16628) from the Ministry of Education, Culture, Sports, Science and Technology, by a Cooperative Research Grant of the Institute for Enzyme Research, Joint Usage/Research Center, Tokushima University, by the Takeda Science Foundation, the Ohya Health Foundation, the Heiwa Nakajima Foundation, the Cell Science Research Foundation, the Mochida Memorial Foundation on Medical and Pharmaceutical Research, the Uehara Memorial Foundation, and the Research Foundation for Microbial Diseases of Osaka University (for MY). This study was also funded by grants-in-aid for Scientific Research (B:17H03913) and Challenging Exploratory Research (19K22940) from the Ministry of Education, Culture, Science, Sports, and Technology (MEXT) of Japan, and by a Livestock Promotional Subsidy from the Japan Racing Association.

References

- Adachi, O., T. Kawai, K. Takeda, M. Matsumoto, H. Tsutsui, M. Sakagami, K. Nakanishi and S. Akira (1998) Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated function. *Immunity*, 9: 143-150.
- Bando, H., Y. Lee, N. Sakaguchi, A. Pradipta, J. S. Ma, S. Tanaka, Y. Cai, J. Liu, J. Shen, Y. Nishikawa, M. Sasai and M. Yamamoto (2018a) Inducible Nitric Oxide Synthase Is a Key Host Factor for *Toxoplasma* GRA15-Dependent Disruption of the Gamma

- Interferon-Induced Antiparasitic Human Response. *MBio*, 9: e01738-18.
- Bando, H., Y. Lee, N. Sakaguchi, A. Pradipta, R. Sakamoto, S. Tanaka, J. S. Ma, M. Sasai and M. Yamamoto (2019) *Toxoplasma* Effector GRA15-Dependent Suppression of IFN-gamma-Induced Antiparasitic Response in Human Neurons. *Frontiers in Cellular and Infection Microbiology*, 9: 140.
- Bando, H., N. Sakaguchi, Y. Lee, A. Pradipta, J. S. Ma, S. Tanaka, D. H. Lai, J. Liu, Z. R. Lun, Y. Nishikawa, M. Sasai and M. Yamamoto (2018b) *Toxoplasma* Effector TgIST Targets Host IDO1 to Antagonize the IFN-gamma-Induced Anti-parasitic Response in Human Cells. *Frontiers in Immunology*, 9: 2073.
- Bando, H., A. Yoshimura, M. Koketsu, A. Soga, Y. Taniguchi, M. Ozaki, M. Suzuki, H. Kanuka and S. Fukumoto (2015) Serological Survey of *Toxoplasma gondii* in Wild Sika Deer in Eastern Hokkaido, Japan. *The Journal of Protozoology Research*, 25: 1-2.
- Batz, M. B., S. Hoffmann and J. G. J. Morris (2012) Ranking the disease burden of 14 pathogens in food sources in the United States using attribution data from outbreak investigations and expert elicitation. *Journal of Food Protection*, 75: 1278-1291.
- Behnke, M. S., A. Khan, J. C. Wootton, J. P. Dubey, K. Tang and L. D. Sibley (2011) Virulence differences in *Toxoplasma* mediated by amplification of a family of polymorphic pseudokinases. *Proceedings of the National Academy of Sciences of the United States of America*, 108: 9631-9636.
- Boothroyd, J. C. (2009) *Toxoplasma gondii*: 25 years and 25 major advances for the field. *International journal for parasitology*, 39: 935-946.
- Courret, N., S. Darce, P. Sonigo, G. Milon, D. Buzoni-Gatel and I. Tardieux (2006) CD11c- and CD11b-expressing mouse leukocytes transport single *Toxoplasma gondii* tachyzoites to the brain. *Blood*, 107: 309-316.
- Dubey, J. P. (2010) *Toxoplasmosis of Animals and Humans*. CRC Press.
- Etheridge, R. D., A. Alagunan, K. Tang, H. J. Lou, B. E. Turk and L. D. Sibley (2014) The *Toxoplasma* pseudokinase ROP5 forms complexes with ROP18 and ROP17 kinases that synergize to control acute virulence in mice. *Cell Host and Microbe*, 15: 537-550.
- Fentress, S. J., M. S. Behnke, I. R. Dunay, M. Mashayekhi, L. M. Rommereim, B. A. Fox, D. J. Bzik, G. A. Taylor, B. E. Turk, C. F. Lichti, R. R. Townsend, W. Qiu, R. Hui, W. L. Beatty and L. D. Sibley (2010) Phosphorylation of immunity-related GTPases by a *Toxoplasma gondii*-secreted kinase promotes macrophage survival and virulence. *Cell Host and Microbe*, 8: 484-495.
- Fisch, D., H. Bando, B. Clough, V. Hornung, M. Yamamoto, A. R. Shenoy and E. M. Frickel (2019) Human GBP1 is a microbe-specific gatekeeper of macrophage apoptosis and pyroptosis. *The EMBO Journal*, 38: e100926.
- Frenkel, J. K. and J. S. Remington (1980) Hepatitis in toxoplasmosis. *The New England journal of medicine*, 302: 178-179.
- Gay, G., L. Braun, M. P. Brenier-Pinchart, J. Vollaie, V. Josserand, R. L. Bertini, A. Varesano, B. Touquet, P. J. De Bock, Y. Coute, I. Tardieux, A. Bougdour and M. A. Hakimi (2016) *Toxoplasma gondii* TgIST co-opts host chromatin repressors dampening STAT1-dependent gene regulation and IFN-gamma-mediated host defenses. *Journal of Experimental Medicine*, 213: 1779-1798.
- Gov, L., A. Karimzadeh, N. Ueno and M. B. Lodoen (2013) Human innate immunity to *Toxoplasma gondii* is mediated by host caspase-1 and ASC and parasite GRA15. *MBio*, 4: e00255-13.
- Gov, L., C. A. Schneider, T. S. Lima, W. Pandori, M. B. Lodoen (2017) NLRP3 and Potassium Efflux Drive Rapid IL-1beta Release from Primary Human Monocytes during *Toxoplasma gondii* Infection. *The Journal of Immunology*, 199: 2855-2864.
- Guzman, E. and M. Montoya (2018) Contributions of Farm Animals to Immunology. *Frontiers in Veterinary Science*, 5: 307.
- Hakimi, M. A., P. Olias and L. D. Sibley (2017) *Toxoplasma* Effectors Targeting Host Signaling and Transcription. *Clinical microbiology reviews*, 30: 615-645.
- Hunter, C. A. and L. D. Sibley (2012) Modulation of innate immunity by *Toxoplasma gondii* virulence effectors. *Nature Reviews Microbiology*, 10: 766-778.
- Jensen, K. D., K. Hu, R. J. Whitmarsh, M. A. Hassan, L. Julien, D. Lu, L. Chen, C. A. Hunter and J. P. Saeij (2013) *Toxoplasma gondii* rhoptry 16 kinase promotes host resistance to oral infection and intestinal inflammation only in the context of the dense granule protein GRA15. *Infection and Immunity*, 81: 2156-2167.
- Jensen, K. D., Y. Wang, E. D. Wojno, A. J. Shastri, K. Hu, L. Cornel, E. Boedec, Y. C. Ong, Y. H. Chien, C. A. Hunter, J. C. Boothroyd and J. P. Saeij (2011) *Toxoplasma* polymorphic effectors determine macrophage polarization and intestinal inflammation. *Cell Host and Microbe*, 9: 472-483.
- Lee, Y., M. Sasai, J. S. Ma, N. Sakaguchi, J. Ohshima, H. Bando, T. Saitoh, S. Akira and M. Yamamoto (2015) p62 Plays a Specific Role in Interferon-gamma-Induced Presentation of a *Toxoplasma* Vacuolar Antigen. *Cell reports*, 13: 223-233.
- Ma, J. S., M. Sasai, J. Ohshima, Y. Lee, H. Bando, K. Takeda and M. Yamamoto (2014) Selective and strain-specific NFAT4 activation by the *Toxoplasma gondii* polymorphic dense granule protein GRA6. *The Journal of experimental medicine*, 211: 2013-2032.
- MacMicking, J. D. (2012) Interferon-inducible effector mechanisms in cell-autonomous immunity. *Nature reviews Immunology*, 12: 367-382.
- Montazeri, M., T. M. Galeh, M. Moosazadeh, S. Sarvi, S. Dodangeh, J. Javidnia, M. Sharif and A. Daryani (2020) The global serological prevalence of *Toxoplasma gondii* in felids during the last five decades (1967-2017): a systematic review and meta-analysis. *Parasites and Vectors*, 13(1): 82.
- Nathan, C. and Q. W. Xie (1994) Nitric oxide synthases: roles, tolls, and controls. *Cell*, 78: 915-918.
- Ohshima, J., M. Sasai, J. Liu, K. Yamashita, J. S. Ma, Y. Lee, H. Bando, J. C. Howard, S. Ebisu, M. Hayashi, K. Takeda, D. M. Standley, E. M. Frickel and M. Yamamoto (2015) RabGDIalpha is a negative regulator of interferon-gamma-inducible GTPase-dependent cell-autonomous immunity to *Toxoplasma gondii*. *Proceedings of the National Academy of Sciences of the United States of America*, 112: E4581-4590.
- Ohshima, J., Y. Lee, M. Sasai, T. Saitoh, J. Su Ma, N. Kamiyama, Y. Matsuura, S. Pann-Ghill, M. Hayashi, S. Ebisu, K. Takeda, S. Akira and M. Yamamoto (2014) Role of mouse and human autophagy proteins in IFN-gamma-induced cell-autonomous responses against *Toxoplasma gondii*. *Journal of Immunology*, 192: 3328-3335.
- Olias, P., R. D. Etheridge, Y. Zhang, M. J. Holtzman and L. D. Sibley (2016) *Toxoplasma* Effector Recruits the Mi-2/NuRD Complex to Repress STAT1 Transcription and Block IFN-gamma-Dependent Gene Expression. *Cell Host and Microbe*, 20: 72-82.
- Platanias, L. C. (2005) Mechanisms of type-I- and type-II-interferon-mediated signaling. *Nature Reviews Immunology*, 5: 375-386.
- Reese, M. L., G. M. Zeiner, J. P. Saeij, J. C. Boothroyd and J. P. Boyle (2011) Polymorphic family of injected pseudokinases is paramount in *Toxoplasma* virulence. *Proceedings of the National Academy of Sciences of the United States of America*, 108: 9625-9630.
- Rosowski, E. E., D. Lu, L. Julien, L. Rodda, R. A. Gaiser, K. D. Jensen and J. P. Saeij (2011) Strain-specific activation of the NF-kappaB pathway by GRA15, a novel *Toxoplasma gondii* dense granule protein. *Journal of Experimental Medicine*, 208: 195-212.
- Rosowski, E. E., Q. P. Nguyen, A. Camejo, E. Spooner and J. P. Saeij (2014) *Toxoplasma gondii* Inhibits gamma interferon (IFN-gamma)- and IFN-beta-induced host cell STAT1 transcriptional activity by increasing the association of STAT1 with DNA. *Infection and Immunity*, 82: 706-719.
- Rosowski, E. E. and J. P. Saeij (2012) *Toxoplasma gondii* clonal strains all inhibit STAT1 transcriptional activity but polymorphic effectors differentially modulate IFN gamma induced gene expression and STAT1 phosphorylation. *PLoS One*, 7: e51448.
- Steinfeldt, T., S. Konen-Waisman, L. Tong, N. Pawlowski, T. Lamkemeyer, L. D. Sibley, J. P. Hunn and J. C. Howard (2010) Phosphorylation of mouse immunity-related GTPase (IRG) resistance proteins is an evasion strategy for virulent *Toxoplasma gondii*. *PLoS biology*, 8: e1000576.
- Stelzer, S., W. Basso, J. Benavides Silván, L. M. Ortega-Mora, P. Maksimov, J. Gethmann, F. J. Conraths and G. Schares (2019) *Toxoplasma gondii* infection and toxoplasmosis in farm animals: Risk factors and economic impact. *Food and Waterborne Parasitology*, 15: e00037.
- Suzuki, Y., M. A. Orellana, R. D. Schreiber and J. S. Remington

- (1988) Interferon-gamma: the major mediator of resistance against *Toxoplasma gondii*. *Science*, 240: 516-518.
- Taylor, G. A., C. G. Feng and A. Sher (2007) Control of IFN-gamma-mediated host resistance to intracellular pathogens by immunity-related GTPases (p47 GTPases). *Microbes and Infection*, 9: 1644-1651.
- Thomas, S. R., D. Mohr and R. Stocker (1994) Nitric oxide inhibits indoleamine 2,3-dioxygenase activity in interferon-gamma primed mononuclear phagocytes. *The Journal of biological chemistry*, 269: 14457-14464.
- Yamamoto, M., M. Okuyama, J. Ma, T. Kimura, N. Kamiyama, H. Saiga, J. Ohshima, M. Sasai, H. Kayama, T. Okamoto, D. C. S. Huang, D. Soldati-Favre, K. Horie, J. Takeda and K. Takeda (2012) A cluster of interferon- γ -inducible p65 GTPases plays a critical role in host defense against *Toxoplasma gondii*. *Immunity*, 37: 302-313.
- Yao, K., J. Fang, Y. L. Yin, Z. M. Feng, Z. R. Tang and G. Wu (2011) Tryptophan metabolism in animals: important roles in nutrition and health. *Frontiers in Bioscience*, 3: 286-297.
- Zhao, Y., D. J. Ferguson, D. C. Wilson, J. C. Howard, L. D. Sibley and G. S. Yap (2009) Virulent *Toxoplasma gondii* evade immunity-related GTPase-mediated parasite vacuole disruption within primed macrophages. *Journal of Immunology*, 182: 3775-3781.